

Radiology Department

Percutaneous Ureteric Stent

Introduction

You have been advised by your hospital consultant that you need to have one or two ureteric stent/s inserted via your kidney/s. This information leaflet tells you about having a Percutaneous Ureteric Stent Insertion. It explains what is involved and what the possible risks are.

This leaflet may not answer all your questions, so if you have any queries or concerns, please do not hesitate to ask the ward staff. If you feel unhappy with any part of your care within the X-ray Department please ask to speak to a senior member of staff.

Please read this leaflet carefully to ensure you are successfully prepared for the examination.

Please contact the X-ray Department if your weight is equal to or more than 205Kg (32 stone) – You may require an alternative examination.

Referral and Consent

If you are having the Percutaneous Ureteric Stent Insertion as a planned procedure, then you should have plenty of time to discuss the situation with your consultant and the radiologist (a specialised X-ray doctor) who will be inserting the percutaneous ureteric stent.

It is possible that you need this procedure done as an emergency, in which case there may be less time for discussion, **but none the less you should have had sufficient explanation before you sign the consent form.** If after discussion with your hospital doctor or radiologist you do not want the procedure carried out, then you can decide against it.

If the radiologist feels that your condition has changed or that your symptoms do not indicate such a procedure is necessary then he/she will explain this to you, communicate with the referring clinician and ask that you return to your referring clinician or ward for review. At all times the radiologist and referring clinician will be acting in your best interests.

Points to Remember

- Please bring any sprays or inhalers that you are taking with you to your appointment.
- We also ask you to leave any valuable possessions on the ward.



Patient Information

- If you are diabetic please note that you are required to have nothing to eat or drink before the examination. Please inform the ward staff so that an alternative medication may be given if necessary.
- If you are taking aspirin or Warfarin please inform the ward staff, as this medication may need to be stopped prior to the procedure.

What is a Percutaneous Ureteric Stent Insertion?

A ureteric stent insertion is a special X-ray procedure, performed under conscious sedation, to relieve the symptoms of an obstructed kidney. The examination will be performed by a radiologist who will be assisted by a radiographer and a radiology nurse.

This procedure is called a percutaneous (meaning through the skin) Ureteric stent (a thin plastic tube that reaches from the kidney down the ureter into the bladder). It does not treat the cause of the blockage but relieves the symptoms.

The urine from a normal kidney drains through a narrow, muscular tube, called the ureter, into the bladder. When the tube becomes blocked, for example by a stone, the kidney can become damaged if left untreated, especially if there is infection present as well. It is likely that you have already had a percutaneous nephrostomy (external drainage tube) inserted into your kidney, to temporarily relieve your symptoms; the ureteric stent will enable urine to bypass the obstruction and drain from your kidney into your bladder. A percutaneous nephrostomy is again left in place at the end of the procedure; however this is normally temporary and removed once it is confirmed that the ureteric stent is draining well. This is usually within a few days following the ureteric stent insertion.

The stent has to be kept in place until the obstruction is relieved. This depends on the cause of obstruction and the nature of its treatment. In the majority of patients, a stent is required for only a short duration, from a few weeks to up to three months without the need to replace it. When the underlying problem is not a kidney stone, the stent can stay in even longer. There are special stents, which may be left in for a much longer time. Your urologist will tell you how long he expects your stent to remain in place.

It is not always easy to predict how complex or how straightforward the procedure will be and thus how long it could take. As a guide, expect to be in the Radiology Department for about 45 minutes to one hour if one stent is inserted; or up to two hours if two stents are inserted into both ureters (bilateral ureteric stent insertion).

Preparation

- Since the procedure will be performed under conscious sedation, **Please do not eat for the 6 hours** prior to your procedure time. You can drink water only during this time but **must stop 2 hours** before your procedure time. This will help reduce the risk of nausea and vomiting following sedation. You should continue with any other usual medications and take it with a small sip of water.
- You will have had some special blood tests to check your blood clotting ability.
- Vitamin K may be required before the procedure if your blood clotting ability is lower than desired.

Patient Information

- You will need a cannula (small plastic tube) inserted into a vein in your arm after you have arrived on the ward or while you are in X-ray Department, to allow administration of intravenous medications, and fluid if needed.
- To reduce the risk of infection, you will be given intravenous antibiotics by a nurse or doctor on the ward prior to going down to X-ray Department.
- The procedure will be explained to you by a clinician and you will be asked to sign a consent form.
- You will need to wear a hospital gown. The Porters will collect you from the ward and bring you to the X-ray department on your bed for the procedure.

During your examination

- After you have arrived in X-ray Department, the procedure will be again explained to you by the radiologist. You will have the chance to ask any further questions that you may have.
- You will be taken into the X-ray room where you will be asked to get onto the X-ray table. You may be asked to lie either on your side, or, if both kidneys are affected, on your tummy.
- You will be attached to monitoring equipment; your vital signs (blood pressure, pulse etc.) will be monitored regularly by a radiology nurse during the procedure.
- You will be given conscious sedation and enough painkillers to help you relax and make you feel sleepy throughout. It is routine for you to be given continuous oxygen through a face mask or nasal cannula during sedation.
- If you do not already have a percutaneous nephrostomy inserted: The radiologist will use the ultrasound machine to decide on the best place for inserting the stent and leaving a temporary percutaneous nephrostomy (external drainage tube and bag) in place. This is usually in your back, just below your ribs.
- Your skin will be cleaned with cold antiseptic solution and sterile drapes will be placed over this area. Then your skin will be numbed with local anaesthetic. When the local anaesthetic is injected, it will sting to start with, but this soon wears off, and the skin and deeper tissues should then feel numb.
- A thin needle will then be inserted into the kidney. When the radiologist is sure that the needle is in the correct position, a guide wire will be passed through the needle into the kidney, down the ureter and into the bladder under X-ray control.
- **If you do already have a percutaneous nephrostomy inserted:** The radiologist will remove the nephrostomy bag and dressing. He/she will then cleanse and prepare your skin as described above, and a guide wire will be passed through the nephrostomy tube into your kidney, down your ureter and into your bladder under X-ray control. The old nephrostomy tube will then be removed.
- Once the guide wire has reached your bladder: The ureteric stent will be passed over the guide wire into your kidney, ureter and bladder. A new catheter (nephrostomy tube) will then be placed into your kidney to act as a temporary external drainage system for a few days following this procedure.
- This nephrostomy tube will then be fixed to the skin surface and attached to a drainage bag. A dressing will be placed on your skin over the catheter.

Patient Information

Risks of the Procedure

The Procedure: X-rays are a type of radiation. We are all exposed to natural background radiation every day of our lives; this comes from the sun, food we eat, and the ground. Exposure to X-rays carries a small risk, but your doctor feels that this risk is outweighed by the benefits of having the test. We will take all safeguards to minimise the amount of X-rays you receive.

As with any procedure or operation complications are possible. We have included the most common risks and complications in this leaflet. The possibility of these complications occurring will vary for each patient and will be discussed with you before you sign the consent form.

Sedation: Sedation can, on rare occasions, cause problems with breathing. If you wish to, this can be discussed with the radiologist at the time of your consultation or before the procedure.

Bleeding: There will be slight bleeding from the kidney. On rare occasions, this may become severe. Your urine will be blood stained but this should start to clear after 24 hours. If it does not start to clear a simple X-ray test (nephrostogram) may be required to find out why.

Infection: Occasionally there may be infection in the kidney. This can usually be treated with antibiotics.

Kidney damage: Very rarely damage may occur to the kidney.

Despite these possible complications, the procedure is normally very safe. At all times during and after the procedure, the staff will be monitoring your responses to this treatment in order to minimize the effects of any complications.

It is possible that you may have an allergic reaction from the contrast agent used. You will be asked about allergies by the radiologist, radiographer or radiology nurse prior to procedure.

Safety

The contrast agent contains iodine and is excreted by the kidneys in your urine. Please inform the radiologist or radiographer:

- If you are allergic to iodine, have any other allergies or suffer from asthma.
- If you have reacted previously to the injection used for kidney X-rays and CT scanning.
- If you are on renal dialysis.

Patients aged 12 – 55 years - could you be pregnant? The risks of radiation are slightly higher for the unborn child so you will be asked to confirm that you are not pregnant before the examination can proceed.

Living with a Ureteric Stent:

Your hospital doctor (urologist) should have discussed the longer term effects of a ureteric stent insertion before you signed the consent form. To summarise, you may experience:

- Increased frequency and urgency to pass urine

Patient Information

- A sensation of incomplete emptying of your bladder
- A small amount of blood in the urine – this can improve with a greater fluid intake
- Slight discomfort or pain in the bladder, kidney, groin or genital area/s

After your examination

- You will be taken back to your ward. Nurses will carry out routine observations, such as taking your pulse, blood pressure and temperature, to make sure that there are no problems. You will stay in bed overnight.
- The nephrostomy tube stays in place attached to a drainage bag. It is important that you take care of this. You should try not to make any sudden movements, for example getting up out of a chair without remembering the bag, and you should make sure that it can move freely with you. The bag will need to be emptied regularly, so that it does not become too heavy, and the nursing staff will want to measure the amount of urine in it each time.
- If you have any problems after the procedure please speak to the staff on the ward

Other Sources of Information

For information about interventional radiology including further information about this procedure you can view the British Society of Interventional Radiology website: www.bsir.org/patients

For information about the effects of X-rays read the NRPB publication: “X-rays how safe are they?” on the Health Protection Agency website: www.hpa.org.uk

Please note that the views expressed in these websites do not necessarily reflect the views of UHCW NHS Trust or the NHS.

The Trust has access to interpreting and translation services. If you need this information in another language or format, please contact the telephone number on your appointment letter and we will do our best to meet your needs.

The Trust operates a smoke free policy.

Document History

Author	Clare Warmington / Yossi Sahin / Amy Lloyd	Reviewed	Sep 2012, Apr 2013, Apr 2015, Dec 2017, July 2018
Department	Radiology Department	Review	July 2020
Contact Tel No	Ext 27161	Version	6
Published	September 2008	Reference	HIC/LFT/740/08