

MRI Aftercare following Intravenous Injection of Contrast Medium, Buscopan, Bowel prep, and Adenosine

Date	Time	Medicine	Volume
		PGD 110 Primovist	
		PGD 111 Gadovist	
		PGD 112 Clariscan/ Dotarem	
		PGD 113 Multihance	
		PGD Rad 1 I.V. Buscopan	
		PGD 115 Mannitol	
		PGD 114 Movi-Prep	
		PGD 89 Klean-prep	
		Regadenoson or Adenosine	
		Abandoned examination	

Name: _____

PID: _____

If you have had I.V Gadolinium (PGD 110,111,112 or 113):

- In the unlikely event that you experience any delayed allergic symptoms (skin rash) within 48 hrs following your injection of contrast, please contact your GP for advice.
- Please also inform the X-ray Department so that details of the reaction can be noted in your MRI record of the examination.
- If bruising is painful and does not appear to improve, please contact your GP for advice.



Patient Information

If you have had I.V. Buscopan:

- The muscles of your eyes may have relaxed, and your vision may be temporarily blurred, this usually lasts up to half an hour. Do not drive until your vision returns to normal.
- If you experience any redness or pain in the eyes over the next 24 hours contact your GP urgently and show him/her this leaflet. This is a very rare occurrence.

If you have had Bowel Preparation (Klean-Prep, Movi-Prep, Mannitol):

- Movi-Prep or Klean-Prep, is a very strong laxative and common possible side effects are nausea, vomiting and thirst. Some bloating or abdominal cramps may occur due to the strong bowel action it produces. You will need easy access to a toilet once you have taken the laxative.
- It is normal to get diarrhoea for 12-24 hours after taking Movi-Prep or Klean-prep. If this occurs, drink plenty of fluid to avoid dehydration.

Patients under the care of a renal specialist should continue to drink from their normal fluid allowance. If you pass a lot of fluid in your bowel motions after taking the laxative, you can relax your fluid restriction to prevent dehydration.

If you are unsure about your fluid allowance in relation to the laxative effect please seek advice from your home care nurse, dialysis nurse or kidney specialist.

If you have had Regadenoson or Adenosine:

- Once you have left the MRI department there should be no side effects related to the administration of Adenosine during your scan.

For further information on any drugs given please visit:

<https://products.mhra.gov.uk>

Abandoned Examinations

If it was not possible to perform your examination today, we will notify the referring clinician with a report which says that we have not performed the imaging which they requested. If you do not hear from your referring team within the next 4 weeks, please contact them to ensure that they have seen the report from radiology.

Patient Information

How to get your results

You will get the results of this examination from the doctor who requested the procedure in one of the following ways:

- Your family doctor (**GP**)

Please make an appointment to see your family doctor once the result is available.

Or

- Your hospital specialist (**OPD consultant**)

If you have not already received an appointment to see him/her in Outpatients, one will be sent to you by his/her clinic co-ordinator.

The Trust has access to interpreting and translation services. If you need this information in another language or format, please contact 024 7696 7117 and we will do our best to meet your needs.

The Trust operates a smoke free policy.

Did we get it right?

We would like you to tell us what you think about our services. This helps us make further improvements and recognise members of staff who provide a good service.

Have your say. Scan the QR code or visit:

www.uhcw.nhs.uk/feedback



Document History

Department:	Radiology
Contact:	27088
Updated:	May 2023
Review:	May 2026
Version:	12
Reference:	HIC/LFT/1482/12