

Day Surgery Unit (Vascular Access Team)


Day Surgery - Everything I need to know about implanted vascular ports

An implanted vascular port is a type of central line which is used for giving intravenous medications and for taking blood samples. They are also known as Ports or porta cath

Implanted ports are small silicone or polyurethane lines which are inserted into a large central vein through a small pocket underneath the skin on the chest or arm.


① Port
(under the skin)

② The line (catheter)
is threaded
through a vein
until the end is
near your heart



Patient Information

The catheter is usually inserted under the skin on your chest. The tip of the line will enter a large central vein just above the heart. Access to the port for administering medicines will be through a small disc which sits in a pocket of skin on the chest. This disc is the size of a 10p coin.



The port will look and feel like a small round bump underneath the skin. You will not be able to see it from outside your body.

The above images were produced by Macmillan Cancer Support and are reused with their permission.

Why should I have an implanted port?

Implanted ports can be used to administer a variety of intravenous treatments. Ports are useful for patients who have small veins and for patients who require long term intravenous therapy.

An implanted port is accessed using a specific type of needle and can be used for:

- The administration of a wide range of intravenous treatments
- The administration of intravenous fluids such as a drip
- Taking blood samples

An implanted port has the following advantages:

Patient Information

- The port is positioned underneath the skin - therefore it cannot be seen from outside your body
- You will be able to swim and bathe with an implanted port in place without worrying about getting it wet
- Implanted ports are for long term treatments
- Once placed and healed, the port does not require any dressings
- The port only needs to be flushed once a month to maintain it

To give treatment through the implanted port, a small specific type of needle called a 'Huber' needle is pushed through the skin directly above the port. A small amount of local anaesthetic gel can be used to numb the skin and make the insertion of the needle into the port less uncomfortable.

Are there any risks?

As with other types of central lines there are risks with an implanted port, however these risks are small. They include:

Infection

Infection of the port can happen at any time while the line is in place. To prevent infection, the line is inserted under sterile conditions. The line will be accessed with strict infection prevention measures to reduce the risk, as infection can occur around the skin pocket of the port or within the bloodstream.

Thrombus or clot

A blood clot may form around the tip of the line. This is often identified when patients have swollen and painful face, neck, or arms or if there is difficulty using the port.

What do I need to before the procedure?

You can eat and drink as normal on the day of surgery.

You need to have your bloods taken within 7 days of your port being inserted.

- If you are on aspirin/clexane or tinzaparin you must not take it the night before surgery.
- If you are on clopidogrel, stop taking it 7 days before surgery.
- If you are on warfarin, ask your consultant when to stop taking it.

What can I expect on the day of the procedure?

- The port will be inserted in theatre usually in the Day Surgery Unit by a surgeon, practitioner, or anaesthetist. A member of staff will be with you throughout. Local anaesthetic will be given to two areas - one in your neck and one in your chest. As it goes in, you will feel a scratch and maybe some stinging.
- You will be covered in sterile drapes to form a tent and an X-ray machine will be over the top of you. A member of staff will be with you to ensure you are comfortable.
- The surgeon/anaesthetist will check that the local anaesthetic has taken effect.
- A small incision is made in your neck to thread the port catheter into your vein. A larger incision, around the size of a 10p coin is then made on your chest to create a pocket of skin to hold the round disc part of the port.
While this is being carried out, you shouldn't feel pain but you may experience feelings of pressure, tugging and popping. If you feel pain, you can tell the member of staff with you and more pain relief can be given.
- Once the port is inserted, you will have a chest X-ray in theatre to confirm the correct position of the port and the tip of the catheter within the vein.
- You will have glue to both wounds as a dressing.
- If you are having chemotherapy soon after your port insertion, you will have a needle in the port and it will be secured with a special dressing.

After the procedure

Once you are on the ward, the local anaesthetic will wear off. You will be able to eat and drink as normal. Your neck and chest may feel tender for the next 24-48 hours after the procedure. Taking your usual painkillers as directed will help to reduce any discomfort.

There will be no dressings required for your port. The cuts will be closed together with stitches under the skin.

You will need to keep this area dry for 5 to 7 days to help with healing.

Care of your implanted port

Your implanted port must be flushed once a month if it is not being used frequently. This will be arranged by the nurses providing your treatment.

You can continue to bathe and shower as normal while you have an implanted port.

How is the implanted port removed?

When your implanted port is no longer required it may be removed.

The removal of the port is a surgical procedure and will take place in theatre under a local anaesthetic.

If you have any questions relating to the insertion, care or removal of an implanted port, please contact the Vascular Access team for advice:

Vascular Access Team

Day Surgery, West wing, Ground Floor

University Hospitals Coventry and Warwickshire NHS Trust

Tel. 024 7696 5900

Patient Information

The Trust has access to interpreting and translation services. If you need this information in another language or format, please contact the Vascular Access Team on 024 7696 5900 and we will do our best to meet your needs.


The Trust operates a smoke free policy.

Did we get it right?

We would like you to tell us what you think about our services. This helps us make further improvements and recognise members of staff who provide a good service.

Have your say. Scan the QR code or visit:

www.uhcw.nhs.uk/feedback



Document History

Department:	Day Surgery Unit
Contact:	26868
Updated:	January 2023
Review:	January 2025
Version:	3
Reference:	HIC/LFT/1870/15