

Department of Nutrition and Dietetics

Allergy - Introducing milk free solids to your baby



This advice is to help you get started with safe, milk free weaning. You should also ask for a referral from your GP or Consultant to see a Dietitian as soon as possible.

Introduction of solids (weaning)

The recommended age for introducing solid foods is by 24 weeks (6 months). This is no different for babies with food allergies or intolerances. It is not advisable to wean before 17 weeks.

One of the most important nutrients to consider for babies with a milk allergy is calcium. This is usually obtained by taking an adequate intake of **breast milk**, or the milk substitute (alternative formula) your GP or Paediatrician has advised for your baby.

Your baby may already be taking one of the alternative formulas such as:

- **Nutramigen 1 LGG or 2 LGG** (2 is suitable from 6 months)
- **Aptamil Pepti and Pepti 2** (2 is suitable from 6 months)
- **Aptamil Pepti Junior**
- **Similac Alimentum**
- **SMA Althéra**
- **Neocate LCP**
- **Puramino**
- **SMA Alfamino**
- **Wysoy (N.B: Soya formulas are not normally recommended under six months of age.)**

Cow's, goat's or sheep's milk are **not suitable** for babies under 12 months, as they do not contain enough vitamins or minerals.

Alternative milks, which are not prescribed, such as oat, soya, almond and coconut milk should not be used as the main source of milk for children under two, but can be used in cooking, and on cereals from 6 months. Rice milk should not be used before 4½ years

How much breast milk/ formula does your baby need?

Breast fed babies or babies on alternative formulas, providing they are gaining weight adequately, will meet their calcium needs. Breast milk contains less calcium than formulas, but it



Patient Information

is absorbed better from breast milk. When your baby starts to eat solids, if their intake of breast milk or formula reduces a little, do not worry. They will get additional calcium from calcium rich foods including; fortified breakfast cereals, milk free yogurts and some breads. Stage 2 formulas, prescribed from 6 months usually have higher calcium content than Stage 1 formulas

If your baby is not taking enough breast milk or formula, they may not be fully meeting their calcium needs, and they may need a calcium supplement. The Dietitian will assess this and will advise on the best way to ensure enough calcium is taken.

Is your baby ready to start solids?

Before you start weaning, check that your baby can sit as upright as possible (slightly inclined if necessary) and they can hold their head well, without being too floppy. They should be showing an interest in others eating and should be starting to bring toys and fingers to their mouths.

Babies are not ready for solids before 17 weeks old

When should I start?

Choose a time when your baby is most awake and most willing to interact. Initially, offer solids at just one feed time. Build up to three meals a day over the course of approximately four to six weeks

What should I give?

Foods such as pureed rice, potatoes, root and green vegetables and fruit are good starting choices if weaning from 17 weeks (see table over page). Use a little of your baby's usual milk (breast or hypoallergenic formula) to mix the food to the desired consistency. If too runny add baby rice or potato to thicken. As babies are starting to wean nearer to six months of age, they can move on to thick, but smooth paste textures after just a few days or weeks

How much is enough?

To start with, give two to three spoons, gradually increasing to a small bowl each meal with a small pudding to follow. Your baby will tell you when they have had enough – try not to force an extra spoon here and there. Their milk feeds may gradually decrease, but they will still need plenty of milk. By seven to eight months most babies will be having three meals daily with main course and pudding at lunch and evening meal. By one year of age they might just have milk in the morning, at bedtime and maybe one in the day. If they drink too much milk after this age, they may not eat enough food.

Allergenic foods

Some foods- nuts, eggs, soya, gluten, wheat, fish, shellfish, sesame and other seeds, are considered to be more likely to cause an allergic reaction.

There is no evidence to support delaying the introduction of these foods later than six months. Soya should not be given before 6 months

Delaying their introduction until over 12 months old may actually increase a child's risk of developing an allergy to that food. Introduction to these allergenic foods should be done one at a time. If there are no adverse reactions observed after 3 days move onto the next food.

Note: If your baby has a fast food allergy (hives, facial swelling, immediate vomiting or difficulty breathing) on their first introduction of these foods, and needs an antihistamine to be given, stop giving it. Further introduction of this food will need to be done under the supervision of an allergy team.

Patient Information

Food Introduction

First weaning (not before 17weeks)	From 6 months (26 weeks)	7 to 9 months	9 to 12 months
Serve food as: Smooth purees	Serve food as: Well-mashed foods	Serve food as: Mashed foods with some lumps and soft finger foods	Serve food as: Mashed, chopped and minced family meals and finger foods
Introduce: Smooth cereals, e.g. baby rice, flaked rice, ground quinoa, cornmeal	Introduce: Gluten containing foods (wheat, barley, rye) – bread and cereals, oats, pasta, couscous	Introduce: Bread, chapatti, cereals, pasta, rice, polenta, noodles, couscous	Introduce: Bread, chapatti, cereals, pasta, rice, polenta, noodles, couscous
Soft cooked pureed/ mashed fruit - bananas, pears, apple, plum, peach, avocado, tomato pureed / sauce, citrus flavours	Soft cooked/ mashed fruit - bananas, pears, apple, plum, peach,, avocado, tomato pureed / sauce, citrus flavours	Raw soft fruit and finger foods e.g. banana, melon, mango, avocado, kiwi, berry fruits Stewed fruit, fresh tomato, citrus fruits	Raw soft fruit and finger foods e.g. banana, melon, mango, avocado, kiwi, berry fruits Stewed fruit, fresh tomato, citrus fruits
Soft cooked root and green vegetables, potatoes, yam, green banana	Mashed root and green vegetables, potatoes, yam, green banana	Raw soft vegetables e.g. cucumber, courgette and lightly cooked vegetables, potatoes, yam, green banana	Raw soft vegetables e.g. cucumber, courgette and lightly cooked vegetables, potatoes, yam, green banana
Pureed pulses/ lentils Meat and poultry (pureed)	Mashed lentils/ pulses Finely chopped/ pureed meat, poultry, fish hummus, smooth nut spreads Yoghurt and cheese alternatives- soya, coconut or oat yoghurt, milk-free cheese, milk free custards/ desserts	Mashed lentils/ pulses Finely chopped/ minced meat, poultry, shellfish & flaked fish, hummus, smooth nut spreads Yoghurt and cheese alternatives - soya, coconut or oat yoghurt, milk-free cheese, milk free custards/ desserts	Mashed lentils/ pulses Chopped/ minced meat, poultry, shellfish, fish, hummus , smooth nut spreads Yoghurt and cheese alternatives - soya, coconut or oat yoghurt, milk-free cheese, milk free custards/ desserts
	British Lion Stamped Egg	British Lion Stamped Egg	British Lion Stamped Egg
	Pureed/mashed boiled, scrambled egg (with breast milk or milk substitute) Smooth peanut butter	Mashed/sliced boiled, scrambled egg (with breast milk or milk substitute) Smooth peanut butter	Chopped boiled egg, scrambled egg (with breast milk or milk substitute) Smooth peanut butter

Manufactured Baby Foods

If you are using manufactured baby foods/jars these may contain milk. You must check labels or ingredient lists carefully. Customer care lines (see on jars for details) can give more information on individual products.

Food Labelling

Every pre-packed food, sold in the UK and European Union (EU) must show clearly on the label if it contains any of 14 major allergens. This includes **milk**. These must be highlighted (in bold, colour or underlined) on food labels within the ingredients list. It is not compulsory for food manufacturers to use a warning statement – always check the ingredients first. However, to help when you go shopping the following list may be useful.

Patient Information

These foods all contain milk and must be avoided:

- **Milk** - Cow's milk - pasteurised, sterilised, UHT, Longlife, semi-skimmed, skimmed, dried, evaporated, condensed.
- **Goat's milk, sheep or ewe's milk.**
- **Cheese and cheese spreads**
- **Butter**
- **Many brands of margarine**
- **Yoghurt and fromage frais**
- **Cream**
- **Ice-cream**
- **Pro-biotic drinks**

It is not always obvious that processed foods contain milk. If a food label lists any of the ingredients below, the **food must be avoided**:

- **Butter / Buttermilk / butter oil**
- **Casein / caseinates / hydrolysed casein**
- **Cheese / cheese curds**
- **Cream and artificial cream**
- **Lactose / lactate / lactitol**
- **Margarine or shortening containing whey**
- **Milk / milk powders / milk solids / non fat milk solids**
- **Whey / hydrolysed whey / whey butter / whey cream / whey protein**
- **Hydrolysed milk protein**
- **Whey syrup sweetener**
- **Yoghurt**

Example daily menu for 1st stage milk free solids

On waking	Breast feed/ special formula
Breakfast	Baby rice/ milk-free cereal with expressed breastmilk or special formula
Mid-morning	Breast feed/ special formula
Lunch	Sieved/ blended/ minced meat/ chicken / fish / lentils Boiled and mashed potato (mashed with breast milk/ special formula) Well-cooked and sieved vegetables Fruit puree
Mid-afternoon	Breast feed/special formula
Evening meal	Custard or pudding made with special formula or fruit puree
Bedtime	Breast feed/special formula

Dental care

Formula milks designed for children with milk allergy contain glucose or corn syrup rather than lactose (milk sugar). These are less kind to teeth. Please pay special attention to your child's dental care.

Always make up the milk to the manufacturer's instructions; do not add more powder to water than is stated.

Patient Information

Once a feed is finished, take away the bottle. Do not use it as a comforter allowing your baby to suck the bottle while going to sleep.

At around six months, introduce a lidded beaker and aim to stop using bottles at one year. As soon as your baby's first tooth comes through, start brushing twice a day with a soft brush.

Vitamin and mineral supplements

The Department of Health recommends that a supplement containing **vitamin D and vitamin A** is given to:

- All breastfed babies from six months of age;
- Breastfed babies from one month old who are at high risk of vitamin D deficiency (i.e. from families with dark pigmented skin where mothers did not take a vitamin D supplement during pregnancy, even if the mother is now taking a vitamin D supplement while breastfeeding);
- Babies under one year who are taking less than 500ml (18floz) of infant formula daily;
- All children from one to five years of age.

If you receive income support, you can obtain **Healthy Start Vitamin drops** from your local baby clinic or health visitor.

Otherwise, you can buy children's vitamins containing vitamins A and D from chemists and supermarkets. Ask your pharmacist for advice if you are unsure which supplement to buy.

Low **calcium** intakes are often a concern for children on milk-free diets, particularly if they are not having enough cow's milk substitute or non-dairy foods that are rich in calcium. If you have concerns, discuss these with your Dietitian who can advise you further.

If your child is avoiding other foods in addition to milk and dairy products, your Dietitian may suggest additional vitamin or mineral supplements.

NOTE:

Breastfeeding mums have higher calcium requirements which can be difficult to meet from diet alone. Unless they are drinking 1000ml of a calcium fortified milk substitute per day or adequate calcium rich foods you may need a supplement. For more information on calcium speak with your Dietitian or see the BDA calcium fact sheet: <https://www.bda.uk.com/foodfacts/Calcium.pdf>. A Vitamin D supplement (10mcg/day) is also recommended.

Further Information and Useful Websites

Allergy UK
FoodMaestro
Spoonguru

This leaflet was produced by Coventry Dietitians, University Hospitals Coventry & Warwickshire NHS Trust. If you have any more questions, please contact the team on **024 7696 6161**.

The Trust has access to interpreting and translation services. If you need this information in another language or format, please contact 024 7696 6161 and we will do our best to meet your needs.

The Trust operates a smoke free policy

Document History – Author: Susan Cooper **Department:** Coventry Dietitians **Contact:** 26161 **Published:** March 2013 **Reviewed:** April 2019 **Review:** April 2021 **Version:** 5.1 **Reference:** HIC/LFT/1578/13