

Upper Limb Unit: Hand Therapy

Finger Extensor Tendons (Early Stage)

You have undergone surgery to repair tendons that straighten your finger(s). It will take approximately 12 weeks for your tendons to be strong enough to return to normal function.

Splint

For the first six weeks the tendon repair is very delicate and can be easily undone. To protect the repair you will be provided with a splint. This splint stops you making a fist which could cause the tendon to re-rupture. You should wear the splint 24 hours a day. Do not use the hand to do any functional tasks as this could also cause the tendon to rupture. You should also avoid hanging or carrying anything from the splint/arm.

You should not drive with your splint on or remove your splint to do so. Driving either with or without your splint puts you at significant risk of rupturing your tendon(s). To drive with your splint on also puts yourself, any passengers and other motorists at risk as you will not have adequate control of the vehicle. You should not drive until at least 8 weeks after your tendon repair.

If your splint is uncomfortable speak to your therapist who can remould or alter your splint. The splint is made out of thermoplastic material which softens when hot so do not expose it to any form of heat.

Wound

Keep your dressing dry and clean at all times. If the dressing gets wet or if you are concerned about your wound contact your GP.



Patient Information

Once your stitches are removed and your wound is fully healed you should start massaging the scar with water-based moisturiser, for example, E45. This is an important part of your treatment as it will help prevent the scar tissue sticking to the tendon and limiting movement of the finger.

Swelling


To minimise swelling keep your hand raised as much as possible.

Exercises

Gentle movement of the tendon also helps to limit scar tissue sticking to the tendon and will begin to gently strengthen the tendon. You should complete only the exercises advised and at the frequency advised. Doing too many exercises may cause inflammation and the formation of extra scar tissue. Doing too few may lead to stiffness.

Remove the finger strap and complete the following exercises every hour.....repetitions of each:

- A. Using your other hand straighten all fingers back fully from the splint.
- B. Gently bend the top two finger joints as far as possible.
- C. Gently lift all your fingers from the splint.



Patient Information

Further Information

Please speak to your therapist regarding any information in this leaflet or contact us:

- Telephone: 024 7696 6013 for Physiotherapy at University Hospitals Coventry and Warwickshire.
- Telephone: 01788 663257 for Physiotherapy at the Hospital of St Cross, Rugby

The Trust has access to interpreting and translation services. If you need this information in another language or format please contact 024 7696 6013 and we will do our best to meet your needs.

The Trust operates a smoke free policy.

Document History

Department:	Upper Limb Unit
Contact:	26016
Updated:	February 2020
Review:	February 2023
Version:	4.1
Reference:	HIC/LFT/1141/10