

Patient Information



Tissue Adhesive/Wound Glue

Parent/Carer Discharge information

This information explains the care of your child's wound which has been closed using tissue adhesive.

What should I do now?

- This special glue does not need to be removed. The glue "scab" will fall off as the wound heals.
- Be aware that there will be a scar, but that it will be a few months before you know what it will look like. The redness may take six months to fade.
- Keep the wound dry. If it becomes wet, gently pat it dry without rubbing. Do not use water-proof dressings.
- Avoid picking or removing any of the glue. It will eventually scab and later come away by itself.
- Do not let your child pick at the wound or dressing.

What medication can I give to my child/adolescent?

Paracetamol and or Ibuprofen (if applicable) to be given for pain, Paracetamol 4-6 hourly last given at..... Ibuprofen 6-8 hourly last given at.....

Return to school/activity

You will need to inform the nursery/school/college that your child has sustained a head injury and may need to refrain from certain activities to ensure full recovery.



Patient Information

When should I seek medical advice?

- Excessive or increasing pain
- Increasing swelling or spreading redness
- Pus or discharge oozing from the wound
- The edges of the wound pull apart

For further advice/information:

- Call NHS 111 (24 hours a day)
- Contact your GP

The Trust has access to interpreting and translation services. If you need this information in another language or format please ask and we will do our best to meet your needs.

The Trust operates a smoke free policy

Document History

Author	Lisa Armour
Department	Children's Emergency
Contact	26918
Published	May 2014
Reviewed	April 2016, April 2018
Review	April 2020
Version	2.1
Reference	HIC/LFT/1755/13