

Endoscopy Unit

Colonoscopy - The procedure explained (second line bowel prep in event of the shortages of standard prep)

The aim of this booklet is to help you make an informed decision about agreeing to the procedure and to help answer any questions that you may have. It also explains the diet you need to follow two days before your appointment.

What is a colonoscopy?

A colonoscopy is an accurate way to look at the lining of your large bowel (colon) to find out if there is any disease or inflammation. It allows us to take tissue samples (biopsies) for analysis by the pathology department if necessary. A colonoscope is used, which is a flexible camera which shows the inside of your bowel on a screen. If a sample needs to be taken, you should not be able to feel it. Photographs are taken and attached to your medical records. If you choose to watch the procedure, please keep in mind that the images on the screen are magnified.

Why do I need a colonoscopy?

A colonoscopy is used to find the cause of your symptoms, help with treatment and to decide if any more treatment or investigation is needed. It is also used for follow-up inspection of previous disease or assessing the clinical importance of an abnormality seen on an X-ray.

Should your condition or symptoms worsen before your



appointment, we advise you to contact your GP immediately.

What are the alternatives?

A barium enema or CT colon are alternatives to a colonoscopy. The advantage of a colonoscopy is that samples can be taken in the same procedure, saving the need for a second examination.

Are there any risks?

A colonoscopy is an invasive procedure, which can lead to complications. Serious complications are extremely rare. The most serious risk is the endoscope damaging your colon during the test. This can cause an infection, bleeding, or a perforation (tear) of the lining of the bowel. If this were to happen, your abdomen (tummy) may become painful and bloated and you may need medicine or surgery to treat the problem.

If a biopsy is taken or a polyp is removed during the test, you may pass a small amount of blood from your anus after the test. This should only happen up to 12–24 hours after the test and is usually no more than a few teaspoons full.

Please note that sometimes the test may need to be stopped or may be incomplete. This can happen if you find the procedure too uncomfortable or if the bowel preparation did not empty your bowel completely. In this case, the test may need to be repeated or we may suggest a different procedure.

What about my medication?

Please continue to take your regular medication as prescribed, unless you are taking any of the following:

- Iron tablets – You must stop **one week before** your appointment.
- Stool bulking agents such as **Fybogel, Regulan, Proctofibre, loperamide (Immodium) or codeine phosphate** – You must stop these **three days before** your appointment.

Patient Information

- **Anticoagulants** – Please contact the unit on 024 7696 6805 if you are taking blood thinning medications such as **warfarin, clopidogrel, rivaroxaban, apixaban, edoxaban, dibigatran, prasugrel or aspirin.**
- **Diabetes** – if your diabetes is controlled with **insulin** or **oral medication**, please make you have received a diabetic information leaflet and that the Endoscopy Unit is aware. This is to make sure you are seen in a timely manner.

Please contact the unit if you have an allergy to Latex.

Preparing for the procedure

You must have a clean and empty lower bowel to ensure the endoscopist has a clear view to complete a full assessment. To do this, you need to follow a strict diet **two days before** your appointment.

Two days before your appointment

Fibre is the indigestible part of cereals, fruit, and vegetables. We ask that you have a low fibre diet on this day.

Food allowed	Food not allowed
All meats (must be lean and tender), fish and eggs	All fruits and vegetables
White breads, pasta, rice, mashed or boiled potatoes (without skin)	Any food containing wholemeal or bran – wholemeal bread, pasta, rice, cereals and biscuits
Yorkshire puddings, pancakes, cream crackers, water biscuits, any pastry containing white flour, crisps	Digestive and coconut biscuits, oat cakes, Ryvita etc
Clear and puree soup	
Tomato and all fruit juice (smooth)	
Boiled sweets, toffees, plain or milk chocolate	

Patient Information

Sponge cake, madeira cake, shortcake	
Ice cream, ice lollies, smooth yoghurt	
All flavours of jelly (except blackcurrant)	
Honey, marmalade, golden and treacle syrup	
Fruit squash, fizzy drinks	
Tea and coffee (small amount of milk)	

If you have specific dietary needs such as coeliac or vegan, please contact the unit on 024 7696 6805.

The day before your procedure

You can have breakfast from foods in the 'allowed' list above.

After this, you must not eat solid, semi-solid or soft food. You should drink as much from the list below as possible:

Fluids Allowed	
Tea and Coffee (small amount of milk)	Tonic/Soda Water, Lemonade
Water	Oxo/Bovril
Smooth Fruit and Tomato Juice	Clear Soup and Broth (no noodles or vegetables)
Fruit Squash	

You may still eat all flavours of jelly (apart from blackcurrant), boiled sweets and clear mints.

Patient Information

If you do start to feel lightheaded, please add sugar or glucose to your drinks.

Do not have thick drinks or soups using flour.

On the day of your procedure

You can continue drinking the allowed fluids until you attend your appointment.

Bowel preparation

You need to take laxatives from the day before your appointment. This will have either been given to you or a prescription sent to you. If you have neither of these then please contact us and we will arrange collection for you from the unit.

The laxatives will make you go to the toilet often so you should stay at home. You may want to apply a barrier cream to protect your bottom.

If you have a morning appointment (before 12 noon) please follow these instructions:

Day before your appointment

- **10:00am** – Take **5 Senna** tablets, with a glass of water.
- **2:00pm** – Dissolve the contents of **one sachet of Picolax/Citrafleet** in **150mls water** and drink over 15 to 20 minutes. The solutions may become hot whilst being mixed. If this happens, wait until it has cooled sufficiently to drink. Please drink a glass (250mls) of allowed fluids every hour to allow the effects of the Picolax/Citrafleet to work. (It may take 2 to 3 hours for the solution to work).
- **6:00pm** - Dissolve the **second sachet of Picolax/Citrafleet** in **150mls water** and drink over 15 to 20 minutes. The solutions may become hot

Patient Information

whilst being mixed. If this happens, wait until it has cooled sufficiently to drink. Please drink a glass (250mls) of allowed fluids every hour to allow the effects of the Picolax/Citrafleet to work. (It may take 2 to 3 hours for the solution to work).

If you have an afternoon appointment (after 12 noon) please follow these instructions:

One day before your appointment

- **2:00pm** - Take **5 Senna** tablets, followed by a glass of water.
- **6:00pm** - Dissolve the contents of **one sachet of Picolax/Citrafleet** in **150mls water** and drink over 15 to 20 minutes. The solutions may become hot whilst being mixed. If this happens, wait until it has cooled sufficiently to drink. Please drink a glass (250mls) of allowed fluids every hour to allow the effects of the Picolax/Citrafleet to work. (It may take 2 to 3 hours for the solution to work).

The day of your appointment

- **07:00am** - Dissolve the **second sachet of Picolax/Citrafleet** in **150mls water** and drink over 15 to 20 minutes. The solutions may become hot whilst being mixed. If this happens, wait until it has cooled sufficiently to drink. Please drink a glass (250mls) of allowed fluids every hour to allow the effects of the Picolax/Citrafleet to work. (It may take 2 to 3 hours for the solution to work)

How long will I be in the Endoscopy Unit?

You should expect to be in the unit for around 3 to 4 hours. The unit may need to prioritise emergency patients from the wards and other departments.

Patient Information

The 3 to 4 hours also depends on how quickly you recover from the procedure, whether you have agreed to have any sedative medication and how busy the department is.

Please note that we do not allow children in the department so please make childcare arrangements or ask for your appointment to be rebooked.

What happens when I arrive?

You will be checked in at reception where the receptionist will check your name, date of birth, address and confirm your arrangements for getting home. You will be given a patient wristband so that staff can safely identify you throughout your stay in Endoscopy Unit.

A member of staff will then take you to an admission room where you will be asked to change into a hospital gown. They will do a brief medical assessment to make sure that you are fit to have the procedure. If you are diabetic, your blood glucose level is checked and recorded. If you take anticoagulant medication, a finger prick test (INR) will also be carried out.

The colonoscopy procedure

You will be taken to the procedure room and introduced to the team, who go through safety checks and answer any last-minute questions. You will then be asked to lie on your left-hand side with your knees bent and feet forward (like you are sitting on a chair). A blood pressure cuff and blood oxygen monitor is placed on your arm and finger to monitor you throughout.

Before the procedure starts, the endoscopist will quickly examine your back passage with a finger, to check for piles.

You may be asked to change position during the procedure; this is to allow the scope to glide through your bowel with ease. There are bends that occur naturally, if needed a member of the team will apply pressure to your abdomen to help the scope through these.

Patient Information

CO₂ gas is used to inflate the bowel; this can result in bloating and cramping. If this happens you will be asked to pass wind. This is perfectly normal and please do not feel embarrassed.

After the procedure

You will be transferred to the recovery area where you will be monitored by a qualified nurse. You should expect to stay in the department unit a safe discharge time is planned. If you are diabetic, your blood glucose will also be checked. Every patient is different. Once you have recovered sufficiently you will be moved into a chair and asked to get dressed. Your lift will be contacted to arrange collection. Before you leave, the nurse or endoscopist may discuss the findings of the procedure with you and give you a discharge leaflet and a copy of your endoscopy report. They will also tell you of any further investigations or treatment you may need.

If you normally suffer from memory problems, a family member or friend can be with you whilst you are being discharged.

You will be given specific information regarding eating and drinking before your discharge. This has been agreed by the endoscopist. Please follow the instructions to prevent complications when you get home.

General points and information

- We aim for you to be seen and treated in a timely manner. However, emergency patients take priority, and we apologise if this delays your appointment. You will be kept updated if this happens.
- You may want to bring a dressing gown and slippers with you.
- The hospital cannot accept responsibility for the loss or damage to personal property during your time on the premises.
- If you have any problems with persistent abdominal pain or bleeding, please contact your GP immediately informing them that you have an appointment with us.
- If you are unable to speak to your GP and your symptoms persist, you must attend Accident and Emergency immediately.
- If you require hospital transport for your appointment, please contact hospital transport direct on 01926 310312.

Patient Information

Further Information

If you need any more information or explanation, please contact the Endoscopy Unit on 024 7696 6805.

This leaflet is based on information produced by the British Society of Gastroenterologists and has been adapted for UHCW NHS Trust by Sakina Lyall, Endoscopy Nurse and Joe Colby Lead Nurse for Gastroenterology.

www.bsg.org.uk

The Trust has access to interpreting and translation services. If you need this information in another language or format, please contact us on 024 7696 6805 and we will do our best to meet your needs.

The Trust operates a smoke free policy.

To give feedback on this leaflet please email feedback@uhcw.nhs.uk

Document History	
Department:	Endoscopy
Contact:	26805
Updated:	August 2022
Review:	August 2024
Version:	12.1
Reference:	HIC/LFT/335/07