

Eye Unit

Posturing following eye surgery: Left cheek to pillow

Following your eye surgery it is important that you posture in the position as advised.

Posturing involves keeping your head in a certain position to ensure that the bubble of gas or oil in your eye remains in the right position, this assists in keeping the retina flat. By keeping your head in a certain position the bubble will float up and towards the problem area in your eye. It is an essential part of your treatment and it is important that you follow the instructions given to you.

You will be required to posture for 50 minutes in every hour. A break of 10 minutes in every hour is allowed. The following information may help you prepare for this part of your treatment which is carried out at home.

It will be explained to you which posturing position is required and some examples are shown overleaf

Practical advice

- Make sure you have suitable chairs/tables/stools/pillows available to ensure you are as comfortable as possible
- Think about what items you may need close by such as: tissues, telephone, radio, drinks (straws may be helpful), extra pillows.
- Ensure you have stocks of food and other essential items. Pre-cooked meals may be useful during this time.
- As a result of the inactivity you may suffer from constipation and/or dehydration so try to have plenty of drinks and fruit/vegetables, and contact your GP if you have concerns or problems.
- Use the 10 minute break for short walks and leg exercises to promote good circulation and relieve pressure on the skin, as well as for putting in eye-drops, comfort breaks, meals, etc.
- Depending on your posturing position, during the 50 minutes you can move about as long as you can keep the correct posturing position.
- Ideally, arrange to have someone with you for the period of time you need to posture, or for someone to provide regular support. Patients that posture may feel isolated or a 'bit down'.



Patient Information

- Sleeping: do your best to stay in the correct posturing position, as far as this is possible. Extra pillows may help you maintain the correct position.
- You may find that the neck muscles ache and become stiff as a result of the posturing. A warm bath or gentle massage may be useful in relieving muscle discomfort. A mild painkiller, such as paracetamol, may also help to relieve muscle stiffness. If you are already taking painkillers for a different condition then continue with these, but **do not take** additional painkillers without medical advice.

What happens to the gas/oil in my eye?

- If gas has been inserted then this will gradually be absorbed and replaced by the natural fluid produced by the eye. Your vision will be poor until the gas bubble is absorbed.
- You will probably be able to see the gas bubble as a line in your vision which slowly moves across your vision as it gradually gets smaller.
- The time taken for the gas bubble to completely disappear is variable but can take up to several weeks.
- **You must not travel in aeroplane until the gas bubble is completely absorbed**, as the gas bubble can expand owing to changes in atmospheric pressure.
- **If you require a general anaesthetic while there is still gas in your eye then you must inform the anaesthetist.**
- If silicone oil is used then this will usually require surgical removal at a later date. You may travel in an aeroplane with silicone oil in the eye as this does not expand.

If you have any queries or concerns about your posturing please contact the ward you were discharged from via the hospital switchboard. Tel. 024 7696 4000.

Details of your posturing position are given are on the following pages.

The Trust has access to interpreting and translation services. If you need this information in another language or format please contact 024 7696 5922 and we will do our best to meet your needs.

The Trust operates a smoke free policy


Document History

Author	Victor Hu / Rosie Johnson/ Mark Hero/ Osama Makhzoum
Department	Ophthalmology
Contact Tel No	25922
Published	April 2012
Reviewed	June 2014, July 2016, October 2018
Review	October 2021
Version	4
Reference No	HIC/LFT/1437/12


Patient Information

Posturing position: LEFT Cheek to Pillow

The Eye



Examples of Posture



Images reproduced with the kind permission of Moorfields Eye Hospital London and Spy Design