

Patient Information

Trauma & Orthopaedics

Shoulder Injury Exercises



To overcome stiffness in your shoulder, do the following exercises.

Remember:

- Remove brace / splint / sling* before exercising. (*Staff to delete as appropriate.)
- Exercise as the pain allows – do not continue if it is too painful!
- Complete five of each exercise.
- As the exercises become easier, build up to completing three times a day.

Should you have any worries or concerns following discharge from hospital, please contact:

Virtual Fracture Clinic Hotline: 024 7696 9237


(Monday to Friday 9am- 4pm)

Whilst the clinic is running, we are unable to take calls so leave a voicemail with your full name and number and you will receive a call back

For emergencies out of these hours please attend A&E.



Patient Information



1. Lean forwards in a steady, supporting position. Swing your arm forward and backwards, slowly like a pendulum.
2. Then swing your arm across your body and out to the side slowly.
3. Complete by swinging your arm in a circular motion.

Exercise images sourced from_Safran, Marc R., MD, Instructions for Sports Medicine Patients, 961-965 by Elsevier

The Trust has access to interpreting and translation services. If you need this Information in another language or format please contact **024 7696 9237** and we will do our best to meet your needs.

The Trust operates a smoke free policy.

Document History

Department: Trauma & Orthopaedics

Contact: 29237

Updated: August 2020

Review: August 2021

Version: 1

Reference: HIC/LFT/2500/20