

This child has the following allergies:

Name:

DOB:






● Watch for signs of ANAPHYLAXIS (life-threatening allergic reaction)

Anaphylaxis may occur without skin symptoms: ALWAYS consider anaphylaxis in someone with known food allergy who has **SUDDEN BREATHING DIFFICULTY**

- | | | |
|--|--|---|
| <p>A AIRWAY</p> <ul style="list-style-type: none"> • Persistent cough • Hoarse voice • Difficulty swallowing • Swollen tongue | <p>B BREATHING</p> <ul style="list-style-type: none"> • Difficult or noisy breathing • Wheeze or persistent cough | <p>C CONSCIOUSNESS</p> <ul style="list-style-type: none"> • Persistent dizziness • Pale or floppy • Suddenly sleepy • Collapse/unconscious |
|--|--|---|

IF ANY ONE (OR MORE) OF THESE SIGNS ABOVE ARE PRESENT:

- 1 Lie child flat with legs raised** (if breathing is difficult, allow child to sit)



- 2 Immediately dial 999** for ambulance and say ANAPHYLAXIS ("ANA-FIL-AX-IS")
- 3** In a school with "spare" back-up adrenaline autoinjectors, **ADMINISTER the SPARE AUTOINJECTOR** if available
- 4** Commence CPR if there are no signs of life
- 5 Stay with child** until ambulance arrives, **do NOT stand child up**
- 6** Phone parent/emergency contact

*** IF IN DOUBT, GIVE ADRENALINE ***

You can dial 999 from any phone, even if there is no credit left on a mobile. Medical observation in hospital is recommended after anaphylaxis. For more information about managing anaphylaxis in schools and 'spare' back-up adrenaline autoinjectors, visit: sparepensinschools.uk

● Mild/moderate reaction:

- Swollen lips, face or eyes
- Itchy/tingling mouth
- Hives or itchy skin rash
- Abdominal pain or vomiting
- Sudden change in behaviour

Action to take:

- Stay with the child, call for help if necessary
- Locate adrenaline autoinjector(s)
- **Give antihistamine:**

(If vomited, can repeat dose)

- Phone parent/emergency contact

Emergency contact details:

1) Name:



2) Name:



Additional instructions:

Parental consent: I hereby authorise school staff to administer the medicines listed on this plan, including a 'spare' back-up adrenaline autoinjector (AAI) if available, in accordance with Department of Health Guidance on the use of AAIs in schools.

Signed:

Print name:

Date:

For more information about managing anaphylaxis in schools and "spare" back-up adrenaline autoinjectors, visit: sparepensinschools.uk

This BSACI Action Plan for Allergic Reactions is for children and young people with mild food allergies, who need to avoid certain allergens. For children at risk of anaphylaxis and who have been prescribed an adrenaline autoinjector device, there are BSACI Action Plans which include instructions for adrenaline autoinjectors. These can be downloaded at bsaci.org

For further information, consult NICE Clinical Guidance CG116 Food allergy in children and young people at guidance.nice.org.uk/CG116

This is a medical document that can only be completed by the child's healthcare professional. It must not be altered without their permission. This document provides medical authorisation for schools to administer a 'spare' adrenaline autoinjector in the event of the above-named child having anaphylaxis (as permitted by the Human Medicines (Amendment) Regulations 2017). The healthcare professional named below confirms that there are no medical contra-indications to the above-named child being administered an adrenaline autoinjector by school staff in an emergency. **This plan has been prepared by:**

Sign & print name:

Hospital/Clinic:



Date: